

Uniform Approximation and Fine Potential Theory*

P. M. GAUTHIER

*Département de mathématiques et de statistique, Université de Montréal,
Montréal, PQ H3C 3J7, Canada*

AND

S. LADOUCEUR

Université Louis Pasteur, Strasbourg, France

Communicated by Allan Pinkus

Received March 20, 1991; accepted October 21, 1991

We give necessary and sufficient conditions for a function defined on a closed subset of \mathbb{R}^N to be the uniform limit of harmonic functions. © 1993 Academic Press, Inc.

Let F be closed subset of \mathbb{R}^N , $N \geq 2$. We denote by $\bar{H}(F)$ the closure in the topology of uniform convergence on F of the space of all harmonic functions on (neighbourhoods of) F .

We shall make use of the fine potential theory for which we refer the reader to [F1, F3, F4].

THEOREM. *Let F be a closed subset of \mathbb{R}^N and u a complex-valued function on F . Then $u \in \bar{H}(F)$ if and only if*

- (1) *u is continuous on F ;*
- (2) *u is finely harmonic on the fine interior of F .*

Proof. The case when F is compact was proved by Debiard and Gaveau [DG] (see also [BH]). For closed sets, the proof follows from the compact case and the localization theorem for harmonic functions on closed sets [GH].

Indeed, suppose that $u \in \bar{H}(F)$. Then, if K is any closed ball, the restric-

* Research supported by NSERC-Canada and FCAR-Quebec.

tion $u|(F \cap K)$ of u to the compact set $F \cap K$ is in $\bar{H}(F \cap K)$. Thus by the Theorem of Debiard and Gaveau, u is continuous on $F \cap K$ and finely harmonic on the fine interior of $F \cap K$. Since this is true for any closed ball K , it follows that u is continuous on F and finely harmonic on the fine interior of F .

Conversely, suppose that u is continuous on F and finely harmonic on the fine interior of F . Again, let K be any closed ball. Since no point of the boundary of K lies in the fine interior of $F \cap K$, it follows that u is continuous on $F \cap K$ and finely harmonic on the fine interior of $F \cap K$. By the Debiard and Gaveau Theorem, $u|(F \cap K) \in \bar{H}(F \cap K)$. Since this is so for every closed ball K , it follows from the localization theorem for harmonic approximation on closed sets [GH, Theorem 2.3.2 and Corollary 2.3.8] that $u \in \bar{H}(F)$.

Remarks. (1) Fine potential theory is usually investigated on domains which admit nonconstant positive superharmonic functions. This would, at first, seem to exclude the plane \mathbb{R}^2 . However, if U is any finely open set in \mathbb{R}^2 , we may define a function to be finely harmonic on U if its restriction to the intersection of U with any ball is finely harmonic.

(2) If Ω is an open set in \mathbb{R}^N and F is a subset of Ω which is closed in the relative topology of Ω , then our theorem (and its proof) still hold. The more general situation where Ω is a Riemannian manifold is currently being considered by Bagby and Blanchet [BB].

(3) An analogous result also holds for approximation by continuous subharmonic functions. For compact sets this is due to Bliedtner and Hansen [BH] (see also [F2]), while for closed sets the result is currently being written [G].

REFERENCES

- [BB] T. BAGBY AND P. BLANCHET, Uniform harmonic approximation on Riemannian manifolds, *J. d'Analyse Math.*, to appear.
- [BH] J. BLIEDTNER AND W. HANSEN, Simplicial cones in potential theory, I, *Invent. Math.* **29** (1975), 83–110; II, *Invent. Math.* **46** (1978), 255–275.
- [DG] A. DEBIARD AND B. GAVEAU, Potentiel fin et algèbres de fonctions analytiques, I, *J. Funct. Anal.* **16** (1974), 289–304; II, *J. Funct. Anal.* **17** (1974), 296–310.
- [F1] B. FUGLEDE, "Finely Harmonic Functions," Lecture Notes in Mathematics, Vol. 289, Springer-Verlag, New York/Berlin, 1972.
- [F2] B. FUGLEDE, Localization in fine potential theory and uniform approximation by subharmonic functions, *J. Funct. Anal.* **49** (1982), 57–72.
- [F3] B. FUGLEDE, Fine potential theory—A survey, *Mitt. Math. Ges. DDR* **2/3** (1986), 3–21.
- [F4] B. FUGLEDE, Fine potential theory, in "Potential Theory Surveys and Problems,"

pp. 81-97, Lecture Notes in Mathematics, Vol. 1344, Springer-Verlag, New York/Berlin, 1988.

[G] P. M. GAUTHIER, Untitled manuscript, in preparation.

[GH] P. M. GAUTHIER AND W. HENGARTNER, "Approximation Qualitative sur des Ensembles Non Bornés," Séminaire de Mathématiques Supérieures, Presses de l'Université de Montréal, 1982.